

HEIGHTS ON THE FINITE PROJECTIVE LINE

MELVYN B. NATHANSON

ABSTRACT. Define the height function $h(a) = \min\{k + (ka \bmod p) : k = 1, 2, \dots, p-1\}$ for $a \in \{0, 1, \dots, p-1\}$. It is proved that the height has peaks at p , $(p+1)/2$, and $(p+1)/3$, that these peaks occur at $a = [p/3]$, $(p-3)/2$, $(p-1)/2$, $[2p/3]$, $p-3$, $p-2$, and $p-1$, and that $h(a) \leq p/3$ for all other values of a .

1. HEIGHTS ON FINITE PROJECTIVE SPACES

Let p be an odd prime and let $\mathbf{F}_p = \mathbf{Z}/p\mathbf{Z}$ and $\mathbf{F}_p^* = \mathbf{F}_p \setminus \{p\mathbf{Z}\}$. For $d \geq 2$, we define an equivalence relation on the set of nonzero d -tuples in $\mathbf{F}_p^d$ as follows: $(a_1 + p\mathbf{Z}, \dots, a_d + p\mathbf{Z}) \sim (b_1 + p\mathbf{Z}, \dots, b_d + p\mathbf{Z})$ if there exists $k \in \{1, 2, \dots, p-1\}$ such that $(b_1 + p\mathbf{Z}, \dots, b_d + p\mathbf{Z}) = (ka_1 + p\mathbf{Z}, \dots, ka_d + p\mathbf{Z})$. We denote the equivalence class of $(a_1 + p\mathbf{Z}, \dots, a_d + p\mathbf{Z})$ by $\langle a_1 + p\mathbf{Z}, \dots, a_d + p\mathbf{Z} \rangle$. The set of equivalence classes is called the $(d-1)$ -dimensional projective space over the field $\mathbf{F}_p$, and denoted $\mathbf{P}^{d-1}(\mathbf{F}_p)$.

For every integer x , we denote by $x \bmod p$ the least nonnegative integer in the congruence class $x + p\mathbf{Z}$. We define the *height* of the point $\mathbf{a} = \langle a_1 + p\mathbf{Z}, \dots, a_d + p\mathbf{Z} \rangle \in \mathbf{P}^{d-1}(\mathbf{F}_p)$ by

$$h_p(\mathbf{a}) = \min \left\{ \sum_{i=1}^d (ka_i \bmod p) : k = 1, \dots, p-1 \right\}.$$

Nathanson and Sullivan [2] introduced this definition of height in connection with a problem of Chudnovsky, Seymour, and Sullivan [1] in graph theory.

The projective line $\mathbf{P}^1(\mathbf{F}_p)$ consists of all equivalence classes of pairs $(a_1 + p\mathbf{Z}, a_2 + p\mathbf{Z})$, where a_1 and a_2 are not both 0 modulo p . If $a_1 \bmod p = 0$, then $\langle a_1 + p\mathbf{Z}, a_2 + p\mathbf{Z} \rangle = \langle p\mathbf{Z}, 1 + p\mathbf{Z} \rangle$ and $h_p(\langle p\mathbf{Z}, 1 + p\mathbf{Z} \rangle) = 1$. If $a_1 \bmod p \neq 0$, then there is an integer a such that $a \equiv a_1^{-1}a_2 \pmod{p}$ and $\langle a_1 + p\mathbf{Z}, a_2 + p\mathbf{Z} \rangle = \langle 1 + p\mathbf{Z}, a + p\mathbf{Z} \rangle$. Thus, for all $\mathbf{a} \in \mathbf{P}^1(\mathbf{F}_p)$, if $\mathbf{a} \neq \langle p\mathbf{Z}, 1 + p\mathbf{Z} \rangle$, then $\mathbf{a} = \langle 1 + p\mathbf{Z}, a + p\mathbf{Z} \rangle$ for some integer $a \in \{0, 1, \dots, p-1\}$. For points on the projective line, we denote the height function by

$$h(a) = h_p(\langle 1, a + p\mathbf{Z} \rangle).$$

In Figure 1 we graph the height function $h(a)$ for $a = 0, 1, \dots, p-2$ for the prime $p = 4409$. Note that $h(p-1) = p$ for all primes p . The graph shows peaks at points $a \approx p/2, p/3, 2p/3, p/4, 3p/4, \dots$ with values $h(a) \approx p/2, p/3, p/3, p/4, p/4, \dots$, and suggests the conjecture that for every positive integer b and all sufficiently large

Date: February 2, 2008.

2000 Mathematics Subject Classification. Primary 11A07, 11G50, 11T99, 05C38.

Key words and phrases. Heights, finite projective line, height spectrum.

The work of M.B.N. was supported in part by grants from the NSA Mathematical Sciences Program and the PSC-CUNY Research Award Program.




FIGURE 1. Heights for the prime 4409

primes p , the range of the height function consists of integers that are equal approximately to $p, p/2, p/3, \dots, p/b$, together with a set of numbers less than p/b . The goal of this paper is to prove this conjecture for $b = 1, 2, 3$.

2. INEQUALITIES AND IDENTITIES

Lemma 1. *Let p be an odd prime number and let m and ℓ be integers such that $1 \leq \ell \leq m \leq p - 1$. If*

$$\frac{(\ell - 1)p}{m} \leq a < \frac{\ell p}{m}$$

then

$$h(a) \leq ma - (\ell - 1)p + m.$$

Proof. Since

$$0 \leq ma - (\ell - 1)p < p,$$

it follows that

$$ma \mod p = ma - (\ell - 1)p.$$

The definition of height implies that

$$h(a) \leq m + (ma \bmod p) = ma - (\ell - 1)p + m.$$

□

Lemma 2. *Let p be an odd prime number and let ℓ and m be integers such that $1 \leq \ell \leq m \leq p - 1$. Let*

$$H_{\ell,m,p}(x) = \ell p - mx - 1 + \frac{mp}{\ell p - mx}$$

for $x < \ell p/m$. If

$$\frac{(\ell - 1)p}{m} < a < \frac{\ell p - 1}{m}$$

then

$$h(a) < H_{\ell,m,p}(a).$$

Let u and v be real numbers such that

$$\frac{(\ell - 1)p}{m} < u \leq \frac{\ell p}{m} - \sqrt{\frac{p}{m}} \leq v < \frac{\ell p - 1}{m}.$$

If $u \leq a \leq v$, then

$$h(a) < \max(H_{\ell,m,p}(u), H_{\ell,m,p}(v)).$$

Proof. We have $1 < \ell p - ma < p$. Dividing the prime number p by $\ell p - ma$, we obtain

$$p = q(\ell p - ma) + r$$

where

$$1 \leq q = \left\lfloor \frac{p}{\ell p - ma} \right\rfloor < \frac{p}{\ell p - ma}$$

and

$$1 \leq r \leq \ell p - ma - 1 < p - 1.$$

Let $k = mq \pmod{p}$. Then $1 \leq k \leq p - 1$ and

$$mqa = p(\ell q - 1) + r$$

and so

$$ka \bmod p = mqa \bmod p = r.$$

It follows that

$$\begin{aligned} h(a) &\leq k + (ka \bmod p) \\ &\leq mq + r \\ &< \ell p - ma - 1 + \frac{mp}{\ell p - ma} \\ &= H_{\ell,m,p}(a). \end{aligned}$$

The function $H_{\ell,m,p}(x)$ is decreasing for $x \leq \ell p/m - \sqrt{p/m}$ and increasing for $\ell p/m - \sqrt{p/m} \leq x < \ell p/m$. It follows that if

$$\frac{(\ell - 1)p}{m} < u \leq \frac{\ell p}{m} - \sqrt{\frac{p}{m}} \leq v < \frac{\ell p - 1}{m}$$

and $u \leq x \leq v$, then $H_{\ell,m,p}(x) \leq \max(H_{\ell,m,p}(u), H_{\ell,m,p}(v))$. This completes the proof. □

Lemma 3. *Let p be an odd prime number and let m and ℓ be integers such that $1 \leq \ell \leq m \leq p-1$. If $m < w \leq \sqrt{mp}$ and*

$$\frac{\ell p}{m} - \frac{p}{w} \leq a \leq \frac{\ell p}{m} - \frac{w}{m}$$

then

$$h(a) < \frac{mp}{w} + w - 1.$$

Proof. We apply Lemma 2 with $u = \ell p/m - p/w$ and $v = (\ell p - w)/m$. If $m < w \leq \sqrt{mp}$, then

$$\frac{(\ell-1)p}{m} < \frac{\ell p}{m} - \frac{p}{w} \leq \frac{\ell p}{m} - \sqrt{\frac{p}{m}} \leq \frac{\ell p}{m} - \frac{w}{m} < \frac{\ell p - 1}{m}.$$

Since the function $H_{\ell,m,p}(w)$ satisfies the functional equation

$$H_{\ell,m,p}\left(\frac{\ell p}{m} - \frac{p}{w}\right) = H_{\ell,m,p}\left(\frac{\ell p}{m} - \frac{w}{m}\right) = \frac{mp}{w} + w - 1$$

it follows that if

$$\frac{\ell p}{m} - \frac{p}{w} \leq a \leq \frac{\ell p}{m} - \frac{w}{m}$$

then

$$h(a) < \max(H_{\ell,m,p}(u), H_{\ell,m,p}(v)) = \frac{mp}{w} + w - 1.$$

This completes the proof. $\square$

Lemma 4. *Let p be an odd prime number and b a positive integer. If $p > (b-1)^2$, then*

$$h(p-b) = \frac{p}{b} + \frac{r(b-1)}{b}$$

where $p \bmod b = r$.

Proof. For each positive integer i , divide ip by b to obtain

$$ip = q_i b + r_i \quad \text{with } 0 \leq r_i \leq b-1.$$

For each i , consider the set of integers k that satisfy the inequality

$$\frac{(i-1)p}{b} < k \leq \frac{ip}{b} = q_i + \frac{r_i}{b}.$$

Then $k \leq q_i$. Every integer $k = 1, 2, \dots, p-1$ satisfies exactly one of these inequalities for some $i \leq b$. If k satisfies the i th inequality, then

$$0 \leq ip - kb < p$$

and

$$k(p-b) \bmod p = ip - kb.$$

It follows that

$$\begin{aligned} k + (k(p-b) \bmod p) &= ip - k(b-1) \\ &\geq ip - q_i(b-1) \\ &= r_i + q_i \\ &= r_i + \frac{ip - r_i}{b} \\ &= \frac{ip}{b} + \frac{r_i(b-1)}{b}. \end{aligned}$$

The inequality $p > (b-1)^2$ implies that $(i+1)p + r_{i+1}(b-1) > ip + r_i(b-1)$ for $i \geq 1$, and so

$$h(p-b) = \min \{k(p-b) \bmod p+k : k=1, \dots, p-1\} \geq \frac{p}{b} + \frac{r_1(b-1)}{b}.$$

Choosing $i=1$ and $k=q_1$ gives $h(p-b) = (p+r_1(b-1))/b$. $\square$

Lemma 5. *Let p be an odd prime number. Then*

$$h\left(\frac{p-1}{2}\right) = \frac{p+1}{2}.$$

If b is an odd integer, $b \geq 3$, and $p > (b-1)^2$, then

$$h\left(\frac{p-b}{2}\right) = \frac{p}{b} + \frac{(b-2)r}{2b}$$

where $p+b \bmod 2b = r$.

Proof. Let $a = (p-1)/2$ and $k \in \{1, \dots, p-1\}$. If $k = 2\ell$ for $\ell \in \{1, \dots, (p-1)/2\}$, then

$$ka \bmod p = \ell(p-1) \bmod p = p-\ell$$

and

$$k + (ka \bmod p) = p + \ell \geq p+1.$$

If $k = 2\ell - 1$ for $\ell \in \{1, \dots, (p-1)/2\}$, then

$$ka \bmod p = \left(\ell(p-1) - \frac{p-1}{2}\right) \bmod p = \frac{p+1}{2} - \ell$$

and

$$k + (ka \bmod p) = \frac{p-1}{2} + \ell \geq \frac{p+1}{2}.$$

It follows that $h((p-1)/2) = (p+1)/2$.

Let $b \geq 3$ and $k \in \{1, 2, \dots, p-1\}$. If k is even, then $k = 2\ell$ for $\ell \in \{1, \dots, (p-1)/2\}$. For each positive integer i , divide ip by b to obtain

$$ip = q_{0,i}b + r_{0,i} \quad \text{with } 0 \leq r_{0,i} \leq b-1.$$

For each i , consider the set of integers ℓ that satisfy the inequality

$$\frac{(i-1)p}{b} < \ell \leq \frac{ip}{b} = q_{0,i} + \frac{r_{0,i}}{b}.$$

Then $\ell \leq q_{0,i}$. Every integer $\ell = 1, 2, \dots, (p-1)/2$ satisfies exactly one of these inequalities for some $i \leq (b+1)/2$. If ℓ satisfies the i th inequality, then

$$0 \leq ip - \ell b < p$$

and

$$k\left(\frac{p-b}{2}\right) \bmod p = \ell(p-b) \bmod p = ip - \ell b.$$

It follows that

$$\begin{aligned}
k + \left(k \left(\frac{p-b}{2} \right) \mod p \right) &= ip - \ell(b-2) \\
&\geq ip - q_{0,i}(b-2) \\
&= r_{0,i} + 2q_{0,i} \\
&= r_{0,i} + \frac{2ip - 2r_{0,i}}{b} \\
&= \frac{2ip}{b} + \frac{(b-2)r_{0,i}}{b}.
\end{aligned}$$

The inequality $p > (b-1)^2$ implies that $2(i+1)p + (b-2)r_{0,i+1} > 2ip + (b-2)r_{0,i}$ for $i \geq 1$, and so, for even integers k , we have

$$(1) \quad \min \{k + (k(p-b) \mod p) : k = 2, 4, \dots, p-1\} \geq \frac{2p}{b} + \frac{(b-2)r_{0,1}}{b}.$$

If k is odd, then $k = 2\ell - 1$ for some $\ell \in \{1, \dots, (p-1)/2\}$. For each positive integer i , divide $(2i-1)p + b$ by $2b$ to obtain

$$(2i-1)p + b = 2bq_{1,i} + r_{1,i} \quad \text{with } 0 \leq r_{1,i} \leq 2b-1.$$

For each i , consider the set of integers ℓ that satisfy the inequality

$$0 \leq ip - \left(\ell b + \frac{p-b}{2} \right) < p,$$

Then $\ell \leq q_{1,i}$. Every integer $\ell = 1, 2, \dots, (p-1)/2$ satisfies exactly one of these inequalities for some $i \leq (b+1)/2$. If ℓ satisfies the i th inequality, then

$$(i-1)p < \ell b + \frac{p-b}{2} \leq ip$$

and

$$k \left(\frac{p-b}{2} \right) \mod p = \ell p - \left(\ell b + \frac{p-b}{2} \right) \mod p = ip - \ell b - \frac{p-b}{2}.$$

It follows that

$$\begin{aligned}
k + \left(k \left(\frac{p-b}{2} \right) \mod p \right) &= ip - \ell(b-2) - \frac{p-b}{2} - 1 \\
&\geq ip - q_{1,i}(b-2) - \frac{p-b}{2} - 1 \\
&= \frac{(2i-1)p}{2} - \left(\frac{(2i-1)p + b - r_{1,i}}{2b} \right) (b-2) + \frac{b}{2} - 1 \\
&= \frac{(2i-1)p}{b} + \frac{(b-2)r_{1,i}}{2b}.
\end{aligned}$$

The inequality $p > (b-1)^2$ implies that

$$2(2i+1)p + (b-2)r_{1,i+1} > 2(2i-1)p + (b-2)r_{1,i}$$

for $i \geq 1$, and so, for odd integers k , we have

$$\min \left\{ k + \left(k \left(\frac{p-b}{2} \right) \mod p \right) : k = 1, 3, \dots, p-2 \right\} \geq \frac{p}{b} + \frac{(b-2)r_{1,1}}{2b}.$$

Choosing $i = 1$ and $\ell = q_{1,1}$, we obtain

$$(2) \quad \min \left\{ k + \left(k \left(\frac{p-b}{2} \right) \mod p \right) : k = 1, 3, \dots, p-2 \right\} = \frac{p}{b} + \frac{(b-2)r_{1,1}}{2b}.$$

Observing that

$$\frac{2p}{b} + \frac{r_{0,1}(b-2)}{b} > \frac{p}{b} + \frac{(b-2)r_{1,1}}{2b}$$

and comparing (1) and (2), we conclude that $h((p-b)/2) = (p+(b-1)r_{1,1})/b$. $\square$

3. PEAKS WITH HEIGHT p/b FOR $b = 1, 2, 3$

Theorem 1. *Let $p \geq 17$ be an odd prime number, and let*

$$A_1 = \{p-1\}$$

and

$$A_2 = \left\{ \left\lfloor \frac{p}{2} \right\rfloor, p-2 \right\}.$$

Then $h(a) = p$ if and only if $a \in A_1$ and $h(a) = (p+1)/2$ if and only if $a \in A_2$. If

$$a \in \{0, 1, 2, \dots, p-1\} \setminus (A_1 \cup A_2)$$

then

$$h(a) \leq \frac{p}{2}$$

Proof. By Lemma 1 with $m = \ell = 1$, we have

$$h(a) \leq a+1 \leq \frac{p-1}{2}$$

for $0 \leq a \leq (p-3)/2$.

By Lemma 1 with $m = \ell = 2$, we have

$$h(a) \leq 2a - p + 2 \leq \frac{p-1}{2}$$

for $(p+1)/2 \leq a \leq (3p-5)/4$.

By Lemma 3 with $m = \ell = 1$ and $w = 3$, we have

$$h(a) < \frac{p}{3} + 2 \leq \frac{p+1}{2}$$

for $2p/3 \leq a \leq p-3$. Note that $2p/3 \leq (3p-5)/4$ for $p \geq 17$. This completes the proof. $\square$

Theorem 2. *Let p be a prime number, and let*

$$A_3 = \left\{ \left\lfloor \frac{p}{3} \right\rfloor, \frac{p-3}{2}, \left\lfloor \frac{2p}{3} \right\rfloor, p-3 \right\}.$$

If $p \equiv 1 \pmod{3}$, then $h(a) = (p+2)/3$ for $a \in A_3$. If $p \equiv 2 \pmod{3}$, then $h(a) = (p+1)/3$ for $a = (p-2)/3$ or $(p-3)/2$ and $h(a) = (p+4)/3$ for $a = (2p-1)/3$ or $p-3$. If

$$a \in \{0, 1, 2, \dots, p-1\} \setminus (A_1 \cup A_2 \cup A_3)$$

then

$$h(a) \leq \frac{p}{3}.$$

Proof. Consider primes $p \equiv 1 \pmod{3}$. Then $p + 3 \equiv 4 \pmod{6}$ and Lemma 5 implies that $h((p-3)/2) = (p+2)/3$. By Lemma 4, $h(p-3) = (p+2)/3$.

We define $q = (p-1)/3$. If $k \in \{1, \dots, p-1\}$, then $k = 3\ell - r$ for $\ell \in \{1, \dots, q\}$ and $r \in \{0, 1, 2\}$. We shall prove that if $a = q$ or $2q$, then $h(a) = \min\{k + (ka \bmod p) : k = 1, \dots, p-1\} = q + 1$. Since $k + (ka \bmod p) \geq k + 1$, it suffices to consider only $k \leq q$ or, equivalently, $\ell \leq (q+r)/3$. We have

$$kq = \frac{(3\ell - r)(p-1)}{3} = \ell(p-1) - rq = \ell p - \ell - rq.$$

Since

$$0 \leq \ell + rq \leq \frac{(3r+1)q+r}{3} \leq \frac{7q+2}{3} < p$$

it follows that $kq \bmod p = p - \ell - rq$ and

$$k + (kq \bmod p) = p + 2\ell - r(q+1).$$

This minimum value of this expression is $q + 1$, and occurs at $\ell = 1, r = 2$, and $k = 1$.

Similarly,

$$k(2q) = \frac{(3\ell - r)(2p-2)}{3} = \ell(2p-2) - 2rq = 2\ell p - 2\ell - 2rq.$$

If $r = 2$, then $p < 2\ell + 4q < 2p$ and $k(2q) \bmod p = 2p - 2\ell - 4q$. Then $k + (k(2q) \bmod p) = 2p + \ell - 4q + 2 = 2q + \ell + 4$. If $r = 0$ or 1 , then $2\ell + 2rq \leq 2\ell + 2q \leq (8q+2)/3 < p$ and $k(2q) \bmod p = p - 2\ell - 2rq$. Then $k + (k(2q) \bmod p) = p + \ell - 2rq - r$ has a minimum value of $q + 1$ at $\ell = 1, r = 1$, and $k = 2$. Thus, $h(a) = q + 1$ if $a \in A_3$.

Next we consider primes $p \equiv 2 \pmod{3}$. Then $p + 3 \equiv 2 \pmod{6}$ and Lemma 5 implies that $h((p-3)/2) = (p+1)/3$. By Lemma 4, $h(p-3) = (p+4)/3$.

Let $q = (p-2)/3$. We shall prove that $h(q) = q + 1$. It suffices to consider only $k + (kq \bmod p)$ for $k = 1, \dots, q$. Let $k = 3\ell - r$ for $r \in \{0, 1, 2\}$. If $k \leq q$, then $\ell \leq (q+r)/3$. We have

$$kq = \frac{(3\ell - r)(p-2)}{3} = \ell p - 2\ell - rq$$

and

$$0 < 2\ell + rq \leq \frac{(2+3r)q+2r}{3} \leq \frac{8q+4}{3} < p.$$

It follows that $kq \bmod p = p - 2\ell - rq$ and so

$$k + (kq \bmod p) = p + \ell - rq - 2 = (3-r)q + \ell.$$

The minimum value of this expression is $q + 1$, and occurs when $\ell = 1, r = 2$, and $k = 1$.

We shall prove that $h(2q+1) = q + 2$. Let $k = 3\ell - r \leq q + 1$, where $r \in \{0, 1, 2\}$ and $1 \leq \ell \leq (q+r+1)/3$. Then

$$k(2q+1) = \frac{(3\ell - r)(2p-1)}{3} = 2\ell p - \ell - 2rq - r.$$

If $r = 2$, then $p < \ell + 4q + 2 = p + q + \ell < 2p$ implies that

$$k(2q+1) \bmod p = 2p - (p + q + \ell) = p - q - \ell$$

and

$$k + k(2q+1) \bmod p = p - q + 2\ell - 2 = 2q + 2\ell \geq 2q + 2.$$

If $r = 0$ or 1 , then

$$0 < \ell + 2rq + r \leq \ell + 2q + 2 < p$$

and

$$k + (k(2q + 1) \bmod p) = (3\ell - r) + (p - \ell - 2rq - r) = p + 2\ell - 2r(q + 1).$$

The minimum value of this expression is $q + 2$ and occurs when $\ell = r = 1$ and $k = 2$.

We have proved that if $a \in A_3$, then $h(a) \in \{(p+1)/3, (p+2)/3, (p+4)/3\}$. We shall prove that if $a \notin A_1 \cup A_2 \cup A_3$, then $h(a) \leq p/3$.

Applying Lemma 1 with $\ell = m = 1$, we obtain $h(a) \leq p/3$ for

$$0 \leq a \leq \frac{p-3}{3}.$$

Applying Lemma 1 with $\ell = 2$ and $m = 3$, we obtain $h(a) \leq p/3$ for

$$\frac{p}{3} \leq a \leq \frac{4p-9}{9}.$$

Applying Lemma 3 with $\ell = 1, m = 2$, and $w = 7$, we have

$$h(a) < \frac{2p}{7} + 6$$

for

$$\frac{5p}{14} \leq a \leq \frac{p-7}{2}.$$

By Lemma 5, we have

$$h\left(\frac{p-5}{2}\right) \leq \frac{p+12}{5}.$$

Since

$$\frac{5p}{14} \leq \frac{4p-9}{9},$$

it follows that if $a \leq (p-5)/2$ and $h(a) > p/3$, then $a = [p/3]$.

Applying Lemma 1 with $\ell = m = 2$, we obtain $h(a) \leq p/3$ for

$$\frac{p}{2} \leq a \leq \frac{2p-4}{3}.$$

Applying Lemma 1 with $\ell = m = 3$, we obtain $h(a) \leq p/3$ for

$$\frac{2p}{3} \leq a \leq \frac{7p-9}{9}.$$

Applying Lemma 3 with $\ell = m = 1$ and $w = 4$, we have

$$h(a) < \frac{p}{4} + 3$$

for

$$\frac{3p}{4} \leq a \leq p-4.$$

Since

$$\frac{3p}{4} \leq \frac{7p-9}{9}$$

it follows that if $p/2 \leq a \leq p-4$ and $h(a) > p/3$, then $a = [2p/3]$. This completes the proof. $\square$

4. PROBLEMS

It is an open problem to describe the spectrum of the height function on projective spaces $\mathbf{P}^{d-1}(\mathbf{F}_p)$ for $d \geq 3$. It would also be interesting to have a reasonable definition of the height function for points in projective space over arbitrary finite fields.

REFERENCES

- [1] M. Chudnovsky, P. Seymour, and B. D. Sullivan, *Cycles in dense digraphs*, arXiv:math.CO/0702147, 2007.
- [2] M. B. Nathanson and B. D. Sullivan, *Heights in finite projective space, and a problem on finite directed graphs*, arXiv: math.NT/0703418, 2007.

DEPARTMENT OF MATHEMATICS, LEHMAN COLLEGE (CUNY), BRONX, NEW YORK 10468
E-mail address: `melvyn.nathanson@lehman.cuny.edu`